11

Likelihood of Physical Contact with Intelligent Extraterrestrial Communities

$$N = R_* f_p n_e f_e f_i f_c^{(N)} L(N)$$

Pessimistic Approach:

$$= 1 \times 10^{-2} \times 3 \times 1 \times 10^{-1} \times 10^{-1} \times L$$

$$= 3 \times 10^{-4} L.$$

Then $L = < 10^3$ yrs $\Longrightarrow N < 0.3$

$L > 10^8$ yrs $\Longrightarrow N > 3 \times 10^4$.

Optimistic Approach:

$$N = 3 \times 10^{-1} \times 3 \times 1 \times 1 \times 0.1 \times L$$

$$= 10^{-1} L.$$

Then $L < 10^3$ yrs $\Longrightarrow N < 10^2$.

$L > 10^8$ yrs $\Longrightarrow N > 10^7$.

For purposes of the following discussion, we adopt $N = 10^6$.

From Sänger (JBIS, 18:273, 1962):

Mass ratio for relativistic rocket with exhaust velocity w:

$$\frac{m_f}{m_i} = \left(\frac{1-\beta}{1+\beta}\right)^{c/2w}$$

where $\beta = v/c$ is the Einstein number (cf. Mach no.).
For a photon rocket, $w = c$. Then

$$\frac{m_f}{m_i} = \left(\frac{1-\beta}{1+\beta}\right)^{1/2}.$$

Let $v = c(1-\varepsilon)$. $\therefore \beta = \frac{v}{c} = 1-\varepsilon$.

$$\therefore \frac{m_f}{m_i} = \left(\frac{1-1+\varepsilon}{1+1-\varepsilon}\right)^{\frac{1}{2}} = \left(\frac{\varepsilon}{2-\varepsilon}\right)^{\frac{1}{2}} \simeq \sqrt{\frac{\varepsilon}{2}}$$

$\therefore \varepsilon = 0.1 \Longrightarrow \frac{m_f}{m_i} \simeq \sqrt{5\times10^{-2}} \simeq 0.2$; $\frac{m_i}{m_f} = 5$

$\varepsilon = 0.01 \Longrightarrow \frac{m_f}{m_i} \simeq \sqrt{5\times10^{-3}} \simeq 0.07$, $\frac{m_i}{m_f} = 14$

$\varepsilon = 0.001 \Longrightarrow \frac{m_f}{m_i} \simeq \sqrt{5\times10^{-4}} \simeq 0.02$; $\frac{m_i}{m_f} = 50$

These are very reasonable mass ratios.

The onboard time under const. acceleration a is

$$t = \frac{2c}{a} \operatorname{arccosh}\left(1 + \frac{aS}{2c^2}\right).$$

where S is the journey distance measured in the rest frame.

Times are doubled to allow for deceleration.

		δ
arccosh	100 000	= 12.2 yr.
arccosh	10^6	= 14.5
arccosh	10^7	= 16.8
arccosh	10^8	= 19.1

11

Consider $a = 980\ cm\ sec^{-2}$ & $S = 1\ l.y. = 9.5\times10^{12}\ km = 9.5\times10^{17}\ cm.$

$$\therefore \frac{aS}{c^2} = \frac{9.5\times10^2\times9.5\times10^{17}}{9\times10^{20}} \simeq 1.$$

$$\therefore t = \frac{3\times10^{10}}{9.8\times10^2}\ \text{arccosh}\ 2 =$$

Now $\cosh 1.32 = 2.$

$$\therefore t = \frac{3\times1.32}{9.8}\times10^8 = 4.0\times10^7\ sec \simeq 1\ yr.$$

$\therefore$ if $a = 1\ g$, it takes 1 yr to reach 1 l.y.

10 l.y. : $\text{arccosh}\ 11 = 3.1$

$$\therefore t = 3.1\times3.1\times10^7 \simeq 3\ yrs.$$

100 l.y. : $\text{arccosh}\ 101 = 5.3$

$$\therefore t = 3.1\times5.3\times10^7 \simeq 5\ yrs$$

1000 l.y. : $\text{arccosh}\ 1001 = 7.6$

$t = 8$ yrs.

10000 l.y. : $\text{arccosh}\ 10001 = 10.0$

$t = 10$ yrs.

For $x > 10$, $\cosh y \simeq \frac{1}{2}e^y.$

$$\therefore y = \text{arccosh}\left[\tfrac{1}{2}e^y\right]$$

Let $\frac{1}{2}e^y = x.$ $\therefore \ln x = \ln\frac{1}{2} + y.$

$$\therefore y = \ln x - \ln\tfrac{1}{2} = \ln 2x$$

$$\therefore \text{arccosh}\ x = \ln 2x$$


S in g's
Interstellar Transit Durations
in Ship Time for a = 1 g
(All times should be doubled to allow
for deceleration)
t in yrs

For intergalactic distances, the eq. reduces to (then accelerate)

$$t \approx \frac{c}{a} \ln [2aS/c^2]$$

The result is that even intergalactic travel becomes reasonable.

However, we choose a time limit on the round trip; say, 10^5 yrs. on the home planet. This restricts travel to within the galaxy.

Assume the gods have lifetimes $\sim 10^2$ yrs. Assume suspended animation in transit. If expedition is to be effective with follow-up (cf Omnes case) several generations of gods required, or suspended animation between visits. [Diff names for Omni encounter → former].
No possibility of sending for another expedition within $\sim 10^2$ yrs, unless base is < 30 l.y. away.

Let each of the N planets in communication phase have q vehicles on continuous patrol. Each vehicle can effect at least one contact per journey. Each vehicle is gone, say, 10^2 yrs.
∴ Taking the lifetime of the communicating phase to be L, the number of contacts effected by each civilization is

$$\sim q \frac{L}{10^2}$$

Total no. contacts by all N's in L yrs is

$$q N L / 10^2$$

∴ No. contacts per yr. is

$qN/10^2$.

With $L = 10^8$ yrs, & $q = 10^2$, & $N = 10^6$, $\exists$

10^6 contacts/yr for all *'s.

10^8 total contacts by each * in L

10^{14} contacts by all *'s in L.

$\therefore$ if contacts are made on a purely random basis, each * should be contacted once each 10^5 yrs if $\exists$ 10^{11} *'s.

We have taken $f_p = 10^{-1}$. The intelligent committee should know which * have planets $\therefore$ on avg. each planetary system contacted randomly once each 10^4 yrs. Where the galactic survey indicates life, & in particular, glimmerings of intelligence, contacts will be more frequent. Man emerged several $\times 10^6$ yrs ago. These events must have been duly noted. $\therefore$ it is quite likely, in the above framework & that the avg. interval between contacts on * during the last 10^4 or 10^5 yrs is $< 10^3$ yrs. $\therefore$ where are they?

Cannes.

Why no subsequent contacts? "In every little boy, we've almost of candy." Hollywood & Portugal

If contact once made, very unlikely that observation is not maintained.

The above treatment tends to depend only inversely on N: because the more unlikely life is, the fewer planetary systems will be investigated provided the survey ships know the parameters involved. There is no distance problem w relativistic spaceflight.

a remarkably straightforward contact history

Oannes is impressive in its directness + in the critical time of the presumed contact. Difficulties w. legends: example in Cosmos + History. Good analogue example in Mead. (European sailing ship contact w. N.W. coast American Indians.) Dragons only an apparent E.T. phylogeny. Bound to be diff. But once, e.g., Tinjanthropus observed could be a selective for galactic community mode like humans to effect contact later. Because of contraction for $\beta \to 1$, it is even possible that Oannes is alive today. Time travel into future.

Acknowledge Barker

Once contact effected, info must go into a central galactic info. repository.

Approach of paper: much recent discussion on methods of contact. Various light, unmanned vehicles. Purpose of the paper is to examine the likelihood of direct contact between galactic civilizations by interstellar spaceflight.

Dragon
ref 1

Communication w. unmanned relativistic rocket almost impossible. Much better to meet at Mead's point contact for pre-communication phase + subsequent. Because of mass limitations, we assume technical assistance: just info.

Pigment; chamber for the gods

(1) questioning of Sumerian mythic (Jacobsen)
(a) cites belief in the gods. Zigguarats. By. and expedition in several cultures over world continuum.

Shepherd (JBIS 11:149, 1952)

$$S = \frac{1}{a^2}\left[\frac{1}{\gamma} - 1\right]; \quad \gamma = \sqrt{1 - \frac{v^2}{c^2}} = \sqrt{1 - \beta^2}$$

Outline:

1. Intro.: Means of communicating. Here study direct physical contact.
2. Number of intelligent communities, N. NAS framework. Acknowledge Drake
3. Interstellar space flight. Techniques, difficulties, and answer argument. Galaxy intercommunicating (Purcell)
4. Consequences. Community organization, central data repository. Frequency of contact. Where are they? Physiological types on other #'s.
5. Possible case of terrestrial contact: the Annedotian Musari.
   (a) Alexander Polyhistor
   (b) Abydenus
   (c) Apollodorus
   (d) Alexander Polyhistor on the Flood (cont.)
6. Connection with early Sumerian history and the subsequent development of civilization.
7. Difficulties in interpreting myths (Elisha, Noah, gods must be human) + other research pass. (Dragons)
8. Future contacts; mode of entry into galactic community


S in light years ⟶
10^7
10^6
10^5
10^4
10^3
10^2
10
6
4
1
7
5
3
2
a = 1g
a = 2g
a = 3g
1 Be V
PROTON
1 Me V
PROTON
1 Ge V
PROTON
1
10^{-1}
10^{-3}
10^{-5}
10^{-7}
10^{-9}
10^{-11}
10^{-13}
10^{-15}
⟵ $\phi = 1 - v/c$

Cab 1019

Bd. of Health CA7-1300 Tom Tonci #38
Boston Health Dept. Haymarket Sq.

10 BeV : $\sim 2\ cm^{-2}\ sec^{-1}$; $\tau \sim 3\times10^{7}\ yrs = 10^{15}\ secs.$
Stopping lifetime (Simpson)

N planets × q ships planet^{-1} yr^{-1} × t yrs any voyage

$N q t$ ships / galaxy

Each ship has velocity $v \simeq c$ + collecting area $A_s\ cm^2$,

$n_i \sim 1\ cm^{-3}$, $n_i v = J\ cm^{-2}\ sec^{-1}$ through collecting area

$\therefore J \sim 3\times10^{10}\ cm^{-2}\ sec^{-1}$.

Attenuated by ratio of areas, A_s/A_g, where A_g = cross-sectional area of galaxy.

$A_s = \pi (3000\ km)^2$

$A_g = \pi (3\times10^{17}\ km)^2$

$\therefore A_s/A_g \simeq (10^3/10^{17})^2 = 10^{-28}$.

1 l.y. = 10^{13} km.
30000 l.y. = 3×10^{17} km

$\therefore 3\times10^{10}\times10^{-28}\times N q t \simeq 2$

$\therefore N q t \sim 10^{18}$

$n_i v A_s$ = no. parts sec^{-1} ship^{-1}

$J_\oplus A_g$ = no. parts sec^{-1} galaxy^{-1}

For $N = 10^6$, $t = 10^3$, $q \sim 10^9$, nope

$$n_i v \times \frac{A_s}{A_g} \times N q t = J_\oplus$$

Should really be $n_i v A_s (N q t)$ no. parts produced by all ships sec^{-1}

$= J_\oplus A_g$ no. parts lost from galaxy sec^{-1}.

$\exists$ N^{cr} ~~parts~~ in volume of galaxy. Lifetime = τ.

$$\therefore \boxed{n_i v \sigma_s N_g t = N/\tau}$$

N: 1 c.r. ~~part~~. cm^{-3} × ~~[illegible]~~ $\pi \times (3\times10^{23})^2 \times 3\times10^{20}$ cm^3

$\simeq 10^{27+46+20} = 10^{68}$ ~~cm~~

$$\begin{array}{r} 68 \\ +1 \\ \hline 27 \end{array} \qquad \begin{array}{r} 15 \\ 10 \\ 16 \\ \hline 41 \end{array}$$

$$\therefore N_g t = \frac{10^{68}}{10^{15}} \times \frac{1}{3\times10^{10} \times \pi \times 4 \times 10^{16}} \sim \frac{1}{30} \times 10^{27} = 3\times10^{25}$$

~~wow~~ yet.

Another difficulty is ~~that~~ 10 BeV ~~proton~~ avg. c.r. energy → voyages only to nearest *'s. ∴ this is a minor contribution to the ~~total~~ galactic c.r. flux.